

## Videnskabelige og faglige artikler

## Fra Steno Diabetes Center 2007



Albrechtsen A, Castella S, Andersen G, Hansen T, Pedersen O, Nielsen R. A Bayesian multilocus association method: Allowing for higher-order interaction in association studies. *Genetics* 2007;176(2):1197-208.

Astrup AS, Nielsen FS, Rossing P, Ali S, Kastrup J, Smidt UM, Parving HH. Predictors of mortality in patients with type 2 diabetes with or without diabetic nephropathy: a follow-up study. *J Hypertens* 2007;25(12):2479-85.

Astrup AS, Tarnow L, Christiansen M, Hansen PR, Parving HH, Rossing P. Pregnancy-associated plasma protein A in a large cohort of Type 1 diabetic patients with and without diabetic nephropathy-a prospective follow-up study. *Diabet Med* 2007;24(12):1381-5.

Berentzen T, Petersen L, Pedersen O, Black E, Astrup A, Sorensen TIA. Long-term effects of leisure time physical activity on risk of insulin resistance and impaired glucose tolerance, allowing for body weight history, in Danish men. *Diabet Med* 2007;24(1):63-72.

Bergholdt R, Stirling ZM, Lage K, Karlberg EO, Olason PI, Aalund M, Nerup J, Brunak S, Workman CT, Pociot F. Integrative analysis for finding genes and networks involved in diabetes and other complex diseases. *Genome Biology* 2007;8(11).

Bergholdt R, Karlsen AE, Hagedorn PH, Aalund M, Nielsen JH, Kruhoffer M, Orntoft T, Wang H, Wollheim CB, Nerup J, Pociot F. Transcriptional profiling of type 1 diabetes genes on chromosome 21 in a rat beta-cell line and human pancreatic islets. *Genes and Immunity* 2007;8(3):232-8.

Bjerregaard P, Jorgensen ME, Borch-Johnsen K. Cardiovascular risk amongst migrant and non-migrant Greenland inuit in a gender perspective. *Scandinavian Journal of Public Health* 2007;35(4):380-6.

Boesgaard TW, Castella SI, Andersen G, Albrechtsen A, Sparso T, Borch-Johnsen K, Jorgensen T, Hansen T, Pedersen O. A-243A -> G polymorphism upstream of the gene encoding GAD65 associates with lower levels of body mass index and glycaemia in a population-based sample of 5857 middle-aged White subjects. *Diabet Med* 2007;24(7):702-6.

Carstensen B. Age-period-cohort models for the Lexis diagram. *Statistics in medicine* 2007 Jul 10;26(15):3018-45.

Chelala C, Duchatelet S, Joffret ML, Bergholdt R, Dubois-Laforgue D, Ghandil P, Pociot F, Caillat-Zucman S, Timsit J, Julier C. PTPN22 R620W functional variant in type 1 diabetes and autoimmunity related traits. *Diabetes* 2007;56(2):522-6.

Dahlman I, Dicker A, Jiao H, Kere J, Blomqvist L, van Harmelen V, Hoffstedt J, Borch-Johnsen K, Jorgensen T, Hansen T, Pedersen O, Laakso M, Arner P. A common haplotype in the G-proteincoupled receptor gene GPR74 is associated with leanness and increased lipolysis. *Am J Hum Genet* 2007;80(6):1115-24.

Eijkelkamp WBA, Zhang ZX, Brenner BM, Cooper ME, Devereux RB, Dahlof B, Ibsen H, Keane WF, Lindholm LH, Olsen MH, Parving HH, Remuzzi G, Shahinfar S, Snapinn SM, Wachtell K, de Zeeuw D. Renal function and risk for cardiovascular events in type 2 diabetic patients with hypertension: the RENAAL and LIFE studies. *J Hypertens* 2007;25(4):871-6.

Eijkelkamp WBA, Zhang ZX, Remuzzi G, Parving HH, Cooper ME, Keane WF, Shahinfar S, Gleim GW, Weir MR, Brenner BM, de Zeeuw D. Albuminuria is a target for renoprotective therapy independent from blood pressure in patients with type 2 diabetic nephropathy: Post hoc analysis from the Reduction of Endpoints in NIDDM with the Angiotensin II Antagonist Losartan (RENAAL) trial. *J Am Soc Nephrol* 2007;18(5):1540-6.

Eising S, Svensson J, Skogstrand K, Nilsson A, Lynch K, Andersen PS, Lernmark A, Hougaard DM, Pociot F, Norgaard-Pedersen B, Nerup J. Type 1 diabetes risk analysis on dried blood spot samples from population-based newborns: design and feasibility of an unselected case-control study. *Paediatr Perinat Epidemiol* 2007;21(6):507-17.

Ellard S, Thomas K, Edghill EL, Owens M, Ambye L, Cropper J, Little J, Strachan M, Stride A, Ersoy B, Eiberg H, Pedersen O, Shepherd MH, Hansen T, Harries LW, Hattersley AT. Partial and whole gene deletion mutations of the GCK and HNF1A genes in maturity-onset diabetes of the young. *Diabetologia* 2007;50(11):2313-7.

Fabrin J, Larsen K, Holstein PE. Arthrodesis with external fixation in the unstable or misaligned Charcot ankle in patients with diabetes mellitus. *Int J Low Extrem Wounds*. 2007 (Jun); 6 (2): 102-107.

Gjesing AP, Andersen G, Albrechtsen A, Glumer C, Borch-Johnsen K, Jorgensen T, Hansen T, Pedersen O. Studies of associations between the Arg389Gly polymorphism of the beta(1)-adrenergic receptor gene (ADRB1) and hypertension and obesity in 7677 Danish white subjects. *Diabet Med* 2007;24(4):392-7.

Gjesing AP, Andersen G, Burgdorf KS, Borch-Johnsen K, Jorgensen T, Hansen T, Pedersen O. Studies of the associations between functional beta(2)-adrenergic receptor variants and obesity, hypertension and type 2 diabetes in 7,808 white subjects. *Diabetologia* 2007;50(3):563-8.

Grarup N, Albrechtsen A, Ek J, Borch-Johnsen K, Jorgensen T, Schmitz O, Hansen T, Pedersen O. Variation in the peroxisome proliferator-activated receptor delta gene in relation to common metabolic traits in 7,495 middle-aged white people. *Diabetologia* 2007;50(6):1201-8.

Grarup N, Andersen G. Gene-environment interactions in the pathogenesis of type 2 diabetes and metabolism. *Current opinion in clinical nutrition and metabolic care* 2007 Jul;10(4):420-6.

Grarup N, Andersen MK, Andreassen CH, Albrechtsen A, Borch-Johnsen K, Jorgensen T, Auwerx J, Schmitz O, Hansen T, Pedersen O. Studies of the common DIO2 Thr92Ala polymorphism and metabolic phenotypes in 7342 Danish white subjects. *J Clin Endocrinol Metab* 2007;92(1):363-6.

Grarup N, Rose CS, Andersson EA, Andersen G, Nielsen AL, Albrechtsen A, Clausen JO, Rasmussen SS, Jorgensen T, Sandbaek A, Lauritzen T, Schmitz O, Hansen T, Pedersen O. Studies of association of variants near the HHEX, CDKN2A/B, and IGF2BP2 genes with type 2 diabetes and impaired insulin release in 10,705 Danish subjects - Validation and extension of genome-wide association studies. *Diabetes* 2007;56(12):3105-11.

Grunnet L, Vielwerth S, Vaag A, Poulsen P. Birth weight is nongenetically associated with glucose intolerance in elderly twins, independent of adult obesity. *J Intern Med* 2007;262(1):96-103.

Gudmundsson J, Sulem P, Steinthorsdottir V, Bergthorsson JT, Thorleifsson G, Manolescu A, Rafnar T, Gudbjartsson D, Agnarsson BA, Baker A, Sigurdsson A, Benediktsdottir KR, Jakobsdottir M, Blondal T, Stacey SN, Helgason A, Gunnarsdottir S, Olafsdottir A, Kristinsson KT, Birgisdottir B, Ghosh S, Thorlacius S, Magnusdottir D, Stefansdottir G, Kristjansson K, Bagger Y, Wilensky RL, Reilly MP, Morris AD, Kimber CH, Adeyemo A, Chen Y, Zhou J, So WY, Tong PCY, Ng MCY, Hansen T, Andersen G, Borch-Johnsen K, Jorgensen T, Tres A, Fuertes F, Ruiz-Echarri M, Asin L, Saez B, van Boven E, Klaver S, Swinkels DW, Aben KK, Graif T, Cashy J, Suarez BK, van Vierssen O, Frigge ML, Ober C, Hofker MH, Wijmenga C, Christiansen C, Rader DJ, Palmer CNA, Rotimi C, Chan JCN, Pedersen O, Sigurdsson G, Benediktsson R, Jonsson E, Einarsson GV, Mayordomo JI, Catalona WJ, Kiemeny LA, Barkardottir RB, Gulcher JR, Thorsteinsdottir U, Kong A, Stefansson K. Two variants on chromosome 17 confer prostate cancer risk, and the one in TCF2 protects against type 2 diabetes. *Nat Genet* 2007;39(8):977-83.

Hadjadj S, Tarnow L, Forsblom C, Kazeem G, Marre M, Groop PH, Parving HH, Cambien F, Tregouet DA, Gut IG, Theva A, Gauguier D, Farrall M, Cox R, Matsuda F, Lathrop M. Association between angiotensin-converting enzyme gene polymorphisms and diabetic nephropathy: Case-control, haplotype, and family-based study in three European populations. *J Am Soc Nephrol* 2007;18(4):1284-91.

Hainerová I, Larsen LH, Holst B, Finková M, Hainer V, Lebl J, Hansen T, Pedersen O. Melanocortin 4 receptor mutations in obese Czech children: studies of prevalence, phenotype development, weight reduction response, and functional analysis. *J Clin Endocrinol Metab* 2007;92: 3689-3696.

Hansen T, Drivsholm T, Urhammer SA, Palacios RT, Volund A, Borch-Johnsen K, Pedersen O. The BIGTT Test - A novel test for simultaneous measurement of pancreatic beta-cell function, insulin sensitivity, and glucose tolerance. *Diabetes Care* 2007;30(2):257-62.

Harman-Boehm I, Sosna T, Lund-Andersen H, Poita M. The eyes in diabetes and diabetes through the eyes. *Diabetes Res Clin Pract* 2007;78:S51-S58.

Haugaard SB, Vaag A, Hoy CE, Madsbad S. Desaturation of skeletal muscle structural and depot lipids in obese individuals during a very-low-calorie diet intervention. *Obesity* 2007;15(1):117-25.

Helgason A, Palsson S, Thorleifsson G, Grant SFA, Emilsson V, Gunnarsdottir S, Adeyemo A, Chen YX, Chen GJ, Reynisdottir I, Benediktsson R, Hinney A, Hansen T, Andersen G, Borch-Johnsen K, Jorgensen T, Schafer H, Faruque M, Doumatey A, Zhou J, Wilensky RL, Reilly MP, Rader DJ, Bagger Y, Christiansen C, Sigurdsson G, Hebebrand J, Pedersen O, Thorsteinsdottir U, Gulcher JR, Kong A, Rotimi C, Stefansson K. Refining the impact of TCF7L2 gene variants on type 2 diabetes and adaptive evolution. *Nat Genet* 2007;39(2):218-25.

Hovind P, Lamberts S, Hop W, Deinum J, Tarnow L, Parving HH, Janssen JAMJ. An IGF-I gene polymorphism modifies the risk of developing persistent microalbuminuria in type 1 diabetes. *European Journal of Endocrinology* 2007;156(1):83-90.

Hughes ML, Carstensen B, Wilkinson JL, Weintraub RG. Angiographic diagnosis, prevalence and outcomes for left ventricular noncompaction in children with congenital cardiac disease. *Cardiol Young*. 2007 (Feb); 17 (1): 56-63.

Hoejberg AS, Hansen D, Svensson J, Johannesen J, Eising S, Birkebaek NH; Danish Registry of Childhood Diabetes. First 10 years with the Danish Registry of Childhood Diabetes. *Ugeskrift for læger*. 2007;28;169(22):2106-8

Jensen CB, Storgaard H, Madsbad S, Richter EA, Vaag AA. Altered skeletal muscle fiber composition and size precede whole-body insulin resistance in young men with low birth weight. *J Clin Endocrinol Metab* 2007;92(4):1530-4.

Jensen DP, Andreasen CH, Andersen MK, Hansen L, Eiberg H, Borch-Johnsen K, Jorgensen T, Hansen T, Pedersen O. The functional Pro129Thr variant of the FAAH gene is not associated with various fat accumulation phenotypes in a population-based cohort of 5,801 whites. *Journal of Molecular Medicine-Jmm* 2007;85(5):445-9.

Jorgensen JOL, Moller L, Krag M, Billestrup N, Christiansen JS. Effects of growth hormone on glucose and fat metabolism in human subjects. *Endocrinol Metab Clin North Am* 2007;36(1):75- 87.

Kamiyama M, Kobayashi M, Araki SI, Iida A, Tsunoda T, Kawai K, Imanishi M, Nomura M, Babazono T, Iwamoto Y, Kashiwagi A, Kaku K, Kawamori R, Ng DPK, Hansen T, Gaede P, Pedersen O, Nakamura Y, Maeda S. Polymorphisms in the 3' UTR in the neurocalcin delta gene affect mRNA stability, and confer susceptibility to diabetic nephropathy. *Hum Genet* 2007;122(3-4):397-407.

Kim WY, Astrup AS, Stuber M, Tarnow L, Falk E, Botnar RM, Simonsen C, Pietraszek L, Hansen PR, Manning WJ, Andersen NT, Parving HH. Subclinical coronary and aortic atherosclerosis detected by magnetic resonance imaging in type 1 diabetes with and without diabetic nephropathy. *Circulation* 2007;115(2):228-35.

Kristensen JK, Drivsholm TB, Carstensen B, Steding-Jensen M, Green A. Validering af metoder til identifikation af erkendt diabetes pa basis af administrative sundhedsregistre. *Ugeskrift for læger* 2007 Apr 30;169(18):1687-92.

Lage K, Karlberg EO, Storling ZM, Olason PI, Pedersen AG, Rigina O, Hinsby AM, Tumer Z, Pociot F, Tommerup N, Moreau Y, Brunak S. A human phenome-interactome network of protein complexes implicated in genetic disorders. *Nat Biotechnol* 2007;25(3):309-16.

Lajer M, Tarnow L, Jorsal A, Parving HH. Polymorphisms in the B-type natriuretic peptide (BNP) gene are associated with NT-proBNP levels but not with diabetic nephropathy or mortality in type 1 diabetic patients. *Nephrology Dialysis Transplantation* 2007;22(11):3235-9.

Larsen CM, Faulenbach M, Vaag A, Volund A, Eshes JA, Seifert B, Mandrup-Poulsen T, Donath MY. Interleukin-1-receptor antagonist in type 2 diabetes mellitus. *N Engl J Med* 2007;356(15):1517-26.

Larsen L, Tonnesen M, Ronn SG, Storling J, Jorgensen S, Mascagni P, Dinarello CA, Billestrup N, Mandrup-Poulsen T. Inhibition of histone deacetylases prevents cytokine-induced toxicity in beta cells. *Diabetologia* 2007;50(4):779-89.

Larsen LH, Rose CS, Sparsø T, Overgaard J, Torekov SS, Grarup N, Jensen DP, Albrechtsen A, Andersen G, Ek J, Glümer C, Borch-Johnsen K, Jorgensen T, Hansen T, Pedersen O. Genetic analysis of the estrogen-related receptor  $\alpha$  and studies of association with obesity and type 2 diabetes. *Int J Obes* 2007;31: 365-370.

Lie BA, Viken MK, Akselsen HE, Flam ST, Pociot F, Nerup J, Kockum I, Cambon-Thomsen A, Thorsby E, Undlien DE. Association analysis in type 1 diabetes of the PRSS16 gene encoding a thymus-specific serine protease. *Hum Immunol* 2007;68(7):592-8.

Ling C, Poulsen P, Simonsson S, Roenn T, Holmkvist J, Almgren P, Hagert P, Nilsson E, Mabey AG, Nilsson P, Vaag A, Groop L. Genetic and epigenetic factors are associated with expression of respiratory chain component NDUFB6 in human skeletal muscle. *J Clin Invest* 2007;117(11):3427-35.

Ling C, Poulsen P, Simonsson S, Rönn T, Holmkvist J, Almgren P, Hagert P, Nilsson E, Mabey A.G, Nilsson P, Vaag A, Groop L. Regulation of skeletal muscle PPAR [delta] mRNA expression in twins. *J Physiol* 584 (Pt 3): 1011-7, 2007.

Ling C, Wegner L, Andersen G, Almgren P, Hansen T, Pedersen O, Groop L, Vaag A, Poulsen P. Impact of the peroxisome proliferator activated receptor-gamma coactivator-1 beta (PGC-1 beta) Ala203Pro polymorphism on in vivo metabolism, PGC-1 beta expression and fibre type composition in human skeletal muscle. *Diabetologia* 2007;50(8):1615-20.

Lund SS, Tarnow L, Stehouwer CDA, Schalkwijk CG, Frandsen M, Smidt UM, Pedersen O, Parving HH, Vaag A. Targeting hyperglycaemia with either metformin or repaglinide in non-obese patients with type 2 diabetes: results from a randomized crossover trial. *Diabetes Obesity & Metabolism* 2007;9(3):394-407.

Mai KS, Sandbaek A, Borch-Johnsen K, Lauritzen T. Are lifestyle changes achieved after participation in a screening programme for Type 2 diabetes? The ADDITION Study, Denmark. *Diabet Med* 2007;24(10):1121-8.

Moller M, Sparre T, Bache N, Roepstorff P, Vorum H. Proteomic analysis of day-night variations in protein levels in the rat pineal gland. *Proteomics* 2007;7(12):2009-18.

Nilsson E, Poulsen P, Sjogren M, Ling C, Ridderstrale M, Groop L, Vaag A. Regulation of skeletal muscle PPAR delta mRNA expression in twins. *Journal of Physiology-London* 2007;584(3):1011-7.

Parikh H, Carlsson E, Chutkow WA, Johansson LE, Storgaard H, Poulsen P, Saxena R, Ladd C, Schulze PC, Mazzini MJ, Jensen CB, Krook A, Bjornholm M, Tornqvist H, Zierath JR, Ridderstrale M, Altschuler D, Lee RT, Vaag A, Groop LC, Mootha VK. TXNIP regulates peripheral glucose metabolism in humans. *Plos Medicine* 2007;4(5):868-79.

Pinterova D, Ek J, Kolostova K, Pruhova S, Novota P, Romzova M, Feigerlova E, Cerna M, Lebl J, Pedersen O, Hansen T. Six novel mutations in the GCK gene in MODY patients. *Clin Genet* 2007;71: 95-96.

Porzio O, Massa O, Cunsolo V, Colombo C, Malaponti M, Bertuzzi F, Hansen T, Johansen A, Pedersen O, Meschi F, Terrinoni A, Melino G, Federici M, Decarlo N, Menicagli M, Campani D, Marchetti P, Ferdaoussi M, Froguel P, Federici G, Vaxillaire M, Barbetti F. Missense mutations in the TGM2 gene encoding transglutaminase 2 are found in patients with early-onset type 2 diabetes. *Hum Mut* 2007;28: 1150.

Poulsen P, Wojtaszewski JFP, Richter EA, Beck-Nielsen H, Vaag A. Low birth weight and zygosity status is associated with defective muscle glycogen and glycogen synthase regulation in elderly twins. *Diabetes* 2007;56(11):2710-4.

Rambrand T, Pociot F, Ronningen KS, Nerup J, Michelsen BK. Genetic markers for glutamic acid decarboxylase do not predict insulin-dependent diabetes mellitus in pairs of affected siblings (vol 99, pg 177, 2007). *Hum Genet* 2007;121(5):653.

Rasmussen SS, Glumer C, Sandbaek A, Lauritzen T, Borch-Johnsen K. Progression from impaired fasting glucose and impaired glucose tolerance to diabetes in a high-risk screening programme in general practice: the ADDITION Study, Denmark. *Diabetologia* 2007;50(2):293-7.

Schloot NC, Hanifi-Moghaddam P, abenhus-Andersen N, Alizadeh BZ, Saha MT, Knip M, Devendra D, Wilkin T, Bonifacio E, Roep BO, Kolb H, Mandrup-Poulsen T. Association of immune mediators at diagnosis of Type 1 diabetes with later clinical remission. *Diabet Med* 2007;24(5):512-20.

Schumann DM, Maedler K, Franklin I, Konrad D, Storling J, Boni-Schnetzler M, Gjinovci A, Kurrer MO, Gauthier BR, Bosco D, Andres A, Berney T, Greter M, Becher B, Chervovsky AV, Halban PA, Mandrup-Poulsen T, Wollheim CB, Donath MY. The Fas pathway is involved in pancreatic beta cell secretory function. *Proc Natl Acad Sci U S A* 2007;104(8):2861-6.

Soja AMB, Zwisler ADO, Frederiksen M, Melchior T, Hommel E, Torp-Pedersen C, Madsen M. Use of intensified comprehensive cardiac rehabilitation to improve risk factor control in patients with type 2 diabetes mellitus or impaired glucose tolerance - the randomized DANish StUdy of impaired glucose metabolism in the settings of cardiac rehabilitation (DANSUK) study. *Am Heart J* 2007;153(4):621-8.

Sparso T, Hussain MS, Andersen G, Hainerova I, Borch-Johnsen K, Jorgensen T, Hansen T, Pedersen O. Relationships between the functional PPAR alpha Leu162Val polymorphism and obesity, type 2 diabetes, dyslipidaemia, and related quantitative traits in studies of 5799 middle-aged white people. *Mol Genet Metab* 2007;90(2):205-9.

Sparso T, Hussain MS, Borch-Johnsen K, Jorgensen T, Madsbad S, Hansen T, Pedersen O, Andersen G. Studies of association of the CASQ1 rs2275703 polymorphism in relation to type 2 diabetes and related quantitative metabolic traits among 7088 Danish whites. *Mol Genet Metab* 2007;92(3):278-82.

Spijkerman AMW, Gall MA, Tarnow L, Twisk JWR, Lauritzen E, Lund-Andersen H, Emeis J, Parving HH, Stehouwer CDA. Endothelial dysfunction and low-grade inflammation and the progression of retinopathy in Type 2 diabetes. *Diabet Med* 2007;24(9):969-76.

Steenkiste A, Valdes AM, Feolo M, Hoffman D, Concannon P, Noble J, Schoch G, Hansen J, Helmberg W, Dorman JS, Thomson G, Pugliese A with the 13th IHWS Type 1 diabetes Component participating investigators. 14th International HLA and immunogenetics workshop: Report on the HLA component of type 1 diabetes. *Tissue Antigens* 69, suppl. 1:214-225, 2007

Steinthorsdottir V, Thorleifsson G, Reynisdottir I, Benediktsson R, Jonsdottir T, Walters GB, Styrkarsdottir U, Gretarsdottir S, Emilsson V, Ghosh S, Baker A, Snorraddottir S, Bjarnason H, Ng MCY, Hansen T, Bagger Y, Wilensky RL, Reilly MP, Adeyemo A, Chen YX, Zhou J, Gudnason V, Chen GJ, Huang HX, Lashley K, Doumatey A, So WY, Ma RCY, Andersen G, Borch-Johnsen K, Jorgensen T, van Vliet-Ostapchouk JV, Hofker MH, Wijmenga C, Christiansen C, Rader DJ, Rotimi C, Gurney M, Chan JCN, Pedersen O, Sigurdsson G, Gulcher JR, Thorsteinsdottir U, Kong A, Stefansson K. A variant in CDKAL1 influences insulin response and risk of type 2 diabetes. *Nat Genet* 2007;39(6):770-5

Storgaard H, Poulsen P, Ling C, Groop L, Vaag AA. Relationships of plasma adiponectin level and adiponectin receptors 1 and 2 gene expression to insulin sensitivity and glucose and fat metabolism in monozygotic and dizygotic twins. *J Clin Endocrinol Metab* 2007;92(7):2835-9.

Svensson J, Carstensen B, Mortensen HB, Borch-Johnsen K. Growth in the first year of life and the risk of type 1 diabetes in a Danish population. *Paediatr Perinat Epidemiol* 2007;21(1):44-8.

Thomson G, Valdes AM, Noble JA, Kockum I, Grote MN, Najman J, Erlich HA, Cucca F, Pugliese A, Steenkiste A, Dorman JS, Caillat-Zucman S, Hermann R, Ilonen J, Lambert AP, Bingley PJ, Gillespie KM, Lernmark A, Sanjeevi CB, Ronningen KS, Undlien DE, Thorsby E, Petrone A, Buzzetti R, Koeleman BPC, Roep BO, Saruhan-Direskeneli G, Uyar FA, Gunoz H, Gorodezky C, Alaez C, Boehm BO, Mlynarski W, Ikegami H, Beririno M, Fasano ME, Dametto E, Israel S, Brautbar C, Santiago-Cortes A, de Llado TF, She JX, Bugawan TL, Rotter JI, Raffel L, Zeidler A, Leyva-Cobian F, Hawkins BR, Chan SH, Castano L, Pociot F, Nerup J. Relative predispositional effects of HLA class II DRB1-DQB1 haplotypes and genotypes on type 1 diabetes: a meta-analysis. *Tissue Antigens* 2007;70(2):110-27

Toft U, Kristoffersen LH, Lau C, Borch-Johnsen K, Jorgensen T. The Dietary Quality Score: validation and association with cardiovascular risk factors: the Inter99 study. *Eur J Clin Nutr* 2007;61(2):270-8.

Tuomilehto J, Jousilahti P, Lindstrom J, Garancini MP, Calori G, Ruotolo G, Bouter LM, Dekker J, Heine RJ, Nijpels G, Stehouwer CDA, Pajak A, Kawalec E, Eliasson M, Stegmayr B, Wareham NJ, Unwin N, Ahmad N, Albert KGMM, Hayes L, Borch-Johnsen K, Qiao Q, Balkau B, Pyorala K. Does diagnosis of the metabolic syndrome detect further men at high risk of cardiovascular death beyond those identified by a conventional cardiovascular risk score? The DECODE Study. *Eur. J. Cardiovasc. Prev. Rehabil.* 2007; 14 (2): 192-199.

Vaiserman AM, Carstensen B, Voitenko VP, Tronko MD, Kravchenko VI, Khalangot MD, Mechova LV. Seasonality of birth in children and young adults (0-29 years) with type 1 diabetes in Ukraine *Diabetologia* 2007;50(1):32-5.

von Huth Smith L, Borch-Johnsen K, Jorgensen T. Commuting physical activity is favourably associated with biological risk factors for cardiovascular disease. *Eur J Epidemiol* 2007;22(11):771-9.

Wagner AM, Cloos P, Bergholdt R, Boissy P, Andersen TL, Henriksen DB, Christiansen C, Christgau S, Pociot F, Nerup J. Post-translational protein modifications in type 1 diabetes: a role for the repair enzyme protein-L-isoaspartate (D-aspartate) O-methyltransferase? *Diabetologia* 2007;50(3):676-81.

Wegner L, Andersen G, Sparso T, Grarup N, Glumer C, Borch-Johnsen K, Jorgensen T, Hansen T, Pedersen O. Common variation in LMNA increases susceptibility to type 2 diabetes and associates with elevated fasting glycemia and estimates of body fat and height in the general population -Studies of 7,495 Danish whites. *Diabetes* 2007;56(3):694-8.

## Oversigtsartikler, lærebøger m.m.

Bergholdt R, Larsen CM, Pociot F, Mandrup-Poulsen T. Type 1 diabetes aetiologi og patogenese (Chapter 6). In: *Diabetes*. Eds.: Hilsted J, Sandahl Christensen J, Borch-Johnsen K. Munksgaard, Denmark, pp 56-65.

Borch-Johnsen K. The metabolic syndrome in a global perspective - The Public Health impact - secondary publication. *Dan Med Bull* 2007;54(2):157-9.

Borch-Johnsen K, Glümer C. Kapitel til Screening-book: Screening for Diabetes. Titel: Screening for Type 2 Diabetes. Screening parameters and techniques, liminations, and opportunities. John Wiley Publishers, 2005.

Borch-Johnsen K. Munksgaard, *Diabetes*, kapitel 2: Diagnose og klassifikation.

Borch-Johnsen K, Pedersen BK. Munksgaard, *Diabetes*, kapitel 8: Forebyggelse af diabetes.

Borch-Johnsen K, Hilsted J. Munksgaard, *Diabetes*, kapitel 31: National database – forskning og kvalitets-sikring.

Fourth Joint Task Force of the European Society of Cardiology and Other Societies on Cardiovascular Disease Prevention in Clinical Practice (Constituted by representatives of nine societies and by invited experts). Authors/Task Force Members, Graham I, Atar D, Borch-Johnsen K et al. European guidelines on cardiovascular disease prevention in clinical practice: executive sum-ary: *Eur Heart J*. 2007 (Oct); 28 (19): 2375-2414.

Gaede P, Pedersen O. Rosiglitazone monotherapy for type 2 diabetes mellitus - too soon to ADOPT? Commentary. *Nature Clinical Practice Endocrinology & Metabolism* 2007;3(6):456-7.

Hansen NW, Bassett B, Borch-Johnsen K. Munksgaard, *Diabetes*, kapitel 4: Screening og opsporing af risikogrupper.

Hansen T. Individualiseret behandling på basis af ny viden om genetikken ved ikke-autoimmun diabetes. *Behandlerbladet* 2007;25: 6-11.

Hilsted J, Borch-Johnsen K, Christiansen JS (editor): *Diabetes*. Munksgaard Danmark.

Larsen CM, Faulenbach M, Vaag A, Vølund Aa, Ehse J, Seifert B, Mandrup-Poulsen T, Donath M (2007) Interleukin-1 Receptor Antagonist behandling af patienter med type 2 diabetes – en sekundærpublikation. *Ugeskrift for læger*. 2007 Nov 5;169(45):3868-71.

Lauritzen T, Borch-Johnsen K, Sandbaek A. Is prevention of Type-2- diabetes feasible and efficient in primary care? A systematic PubMed review. *Primary Care Diabetes* 2007 (Feb); 1 (1): 5-11.

Mandrup-Poulsen TR. Glutaminsyre-decarboxylase-antistoffer og diabetes. *Ugeskrift for læger* 2007 Jan 22;169(4):300-4.

Nerup J, Pociot F. Onset of type 1 diabetes: A dynamic instability. *Mathematic modelling of type 1 diabetes pathogenesis - where did it take us? Commentaries on Perspectives in Diabetes* 3 (1998–2002): 165-166, 2007

Pedersen O. Reducing cardiovascular disease risk in type 2 diabetes: mission possible. *Medicographia* 29: 266-272.

Poulsen P, Esteller M, Vaag A, Fraga MF. The epigenetic basis of twin discordance in age-related diseases. *Pediatr Res* 2007;61(5):38R-42R.

Ronn SG, Billestrup N, Mandrup-Poulsen T. Diabetes and suppressors of cytokine signaling proteins. *Diabetes* 2007;56(2):541-8.

Rossing K. Progression and remission of nephropathy in type 2 diabetes: new strategies of treatment and monitoring. *Dan Med Bull* 2007;54(2):79-98.

Sonne FP, Scheede-Bergdahl C, Olsen DB, Hojbjerg R, Alibegovic A, Nielsen NB, Stallknech B, Helge JW, Vaag A, Dela F. Effects of physical training on endothelial function and limb blood flow in type 2 diabetes. *Applied Physiology Nutrition and Metabolism-Physiologie Appliquee Nutrition et Metabolisme* 2007;32(5):936-41.

Vaag A, Lund SS. Non-obese patients with type 2 diabetes and prediabetic subjects: distinct phenotypes requiring special diabetes treatment and (or) prevention? *Applied Physiology Nutrition and Metabolism-Physiologie Appliquee Nutrition et Metabolisme* 2007;32(5):912-20.

Vaag A, Poulsen P. Twins in metabolic and diabetes research: what do they tell us? *Current opinion in clinical nutrition and metabolic care* 2007 Sep;10(5):591-6.

Vaag A, Schmitz O. Perorale antidiabetika. *Munksgaard Danmark 'Diabetes'* 13:13 a, 2006.

## Ph.d.- afhandlinger

Froboese H. Effects of SOCS-3 on beta-cell toxicity in diabetes.

## Disputatser

Rossing K. Progression and remission of nephropathy in type 2-diabetes. Disputats. [Københavns Universitet. Det Sundhedsvidenskabelige Fakultet] 2007.

## Specialeafhandlinger

Bach BH. Vurdering af eksisterende metoder til måling af fysisk aktivitet i populationsstudier. Københavns Universitet.

Bang-Bertelsen C. Characterization of crb-1 in T1D pathogenesis.

Bork-Jensen J. Variation in genome-wide gene expression in monozygotic twins discordant for glucose tolerance.

Burgdorf KS. Studies of genetic risk factors in KCNMB1, AHSB, and KLF7: candidate genes in the pathogenesis of type 2 diabetes, obesity, dyslipidaemia, and hypertension.

Hussain M. Studies of genetic risk factors in the pathogenesis of type 2 diabetes, obesity, and dyslipidemia: experimental focus on the candidate genes PPARA, FOXO1A, and TCF7L2.

Jacobsen AS. Effect of short term high fat overfeeding on XPHOS gene expression in skeletal muscle of normal- and low birth weight subjects.

Jakobsen MLB. The effects of SOCS-3 on cytokine-induced chemokine expression in pancreatic  $\beta$ -cells.

Lilleoere SK. The effect of birth weight on 24-hour energy expenditure, respiratory quotient and substrate oxidation during a weight-maintenance diet and a short-term high-fat overfeeding diet.

Prause MC. The role of TNF receptor-associated factor 2 in IL-1 $\beta$  + IFN $\gamma$ -induced, endoplasmic reticulum stress-mediated pancreatic  $\beta$ -cell death.

Rask MB. Insulin resistant individuals without metabolic abnormalities and insulin sensitive individuals with metabolic abnormalities – How to explain the paradox? Københavns Universitet.

Voldby LA. Socioeconomic status and screening for Type 2 Diabetes Mellitus – Can socioeconomic factors identify individuals with undiagnosed diabetes? Københavns Universitet. subjects and matched controls.